



HOW TO USE: Read the medication guide provided by your pharmacist before you start taking this medication and each time you get a refill. If you have any questions, consult your doctor or pharmacist.

Your doctor should tell you how long before the procedure you should take this product. Follow your doctors' directions closely. If you are uncertain about any of the information, consult your doctor or pharmacist.

Follow any special dietary instructions given by your doctor. A period of not eating is usually recommended before you start this medication and, lasting until after the colonoscopy. Only clear liquids (such as water, flavored water, lemonade with no pulp, ginger ale, apple juice) are allowed during this time. **Do not drink any liquids colored purple or red.**

The evening before colonoscopy, take 4 tablets with at least 8 ounces (240 ml) of clear liquid by mouth every 15 minutes or as directed by your doctor. Repeat this dose and the liquids every 15 minutes until you have taken 5 doses (20 tablets). The next day, start taking the medication 3-5 hours before the colonoscopy. Take 4 tablets every 15 minutes with liquids as directed until you have taken 3 doses (12 tablets).

Abu Hurairah, MD | Anne Joseph, APRN | Meghana Vellanki, MD
Treece Lewis, APRN | Vishwas Vanar, MD

2100 Ocoee Apopka Rd., Ste. 110 Apopka, FL. 32703
2000 Fowlers Grove Blvd., 3rd Floor Winter Garden, FL. 34787
1919 E. Highway 50, Ste. 201 Clermont, FL. 34711
Phone: 407-609-7395 | Fax: 407-609-7297